

FISH & RICHARDSON P.C.

A

45 Rockefeller Plaza
Suite 2800
New York, New York
10111

Telephone
212 765-5070

Facsimile
212 258-2291

Web Site
www.fr.com

66/09/99
Jc583 U.S. PTO
Frederick P. Fish
1855-1930
W.K. Richardson
1859-1951

September 9, 1999

Attorney Docket No.: 11112/002001

Box Patent Application

Assistant Commissioner for Patents
Washington, DC 20231

Presented for filing is a new patent application claiming priority from a provisional patent application of:

Applicant: Steven P. Nolan and Jinkun Huang
Title: CATALYST COMPLEX WITH CARBENE LIGAND

Enclosed are the following papers, including those required to receive a filing date under 37 CFR §1.53(b):

	<u>Pages</u>
Specification	29
Claims	10
Abstract	1
Signed Declaration	[To Be Filed At A Later Date]
Drawing(s)	6

Enclosures:

- Small entity statement. This application is entitled to small entity status.
- Postcard.

Under 35 USC §119(e)(1), this application claims the benefit of prior U.S. Provisional Application Serial Nos. 60/009,722, filed September 10, 1998, and 60/115,358, filed January 8, 1999.

"EXPRESS MAIL" Mailing Label Number E137209A18905

Date of Deposit September 9, 1999
I hereby certify under 37 CFR 1.10 that this correspondence is being deposited with the United States Postal Service as "Express Mail Post Office To Addressee" with sufficient postage on the date indicated above and is addressed to the Assistant Commissioner for Patents, Washington, D.C. 20231.

Francisco Robles
Francisco Robles

BOSTON
DELAWARE
NEW YORK
SILICON VALLEY
SOUTHERN CALIFORNIA
TWIN CITIES
WASHINGTON, DC

Jc135 U.S. PTO
66/09/99
09/37/869

FISH & RICHARDSON P.C.

September 9, 1999

Page 2

Basic filing fee	380.00
Total claims in excess of 20 times \$9.00	54.00
Independent claims in excess of 3 times \$39.00	78.00
Fee for multiple dependent claims	0.00
Total filing fee:	\$ 512.00

A check for the filing fee is enclosed. Please apply any other required fees or any credits to deposit account 06-1050, referencing the attorney docket number shown above.


If this application is found to be incomplete, or if a telephone conference would otherwise be helpful, please call the undersigned at 212/765-5070.

Kindly acknowledge receipt of this application by returning the enclosed postcard.

Please send all correspondence to:

Scott C. Harris
Fish & Richardson P.C.
4225 Executive Square, Ste. 1400
La Jolla, CA 92037

Respectfully submitted,


George E. Heibel
Reg. No. 42,648
Enclosures

66000 " 63222250

Applicant or Patentee: Steven P. Nolan and Jinkun Huang
 Serial or Patent No.:
 Filed or Issued:
 For: CATALYST COMPLEX WITH CARBENE LIGAND

VERIFIED STATEMENT (DECLARATION) CLAIMING SMALL ENTITY STATUS
 (37 CFR 1.9(c) and 1.27(c)) - NONPROFIT ORGANIZATION

I hereby declare that I am

- ☐ the owner of the small business concern identified below:
☒ an official of the small business concern empowered to act on behalf of the concern identified below:

Name of Small Business Concern: University of New Orleans Foundation

Address of Small Business Concern: Office of Research, Administration Building 208, New Orleans, LA 70148

I hereby declare that the above identified small business concern qualifies as a small business concern as defined in 37 CFR 1.21.12, and reproduced in 37 CFR 1.9(d), for purposes of paying reduced fees to the United States Patent and Trademark Office, in that the number of employees of the concern, including those of its affiliates, does not exceed 500 persons. For purposes of this statement, (1) the number of employees of the business concern is the average over the previous fiscal year of the concern of the persons employed on a full-time, part-time or temporary basis during each of the pay periods of the fiscal year, and (2) concerns are affiliates of each other when either, directly or indirectly, one concern controls or has the power to control the other, or a third party or parties controls or has the power to control both.

I hereby declare that rights under contract or law have been conveyed to and remain with the small business concern identified above with regard to the invention, entitled Office of Research by Inventor(s) Administration Building 208 described in

- ☒ the specification filed herewith.
☐ application serial no. , filed .
☐ patent no. , issued .

If the rights held by the above identified small business concern are not exclusive, each individual, concern or organization having rights to the invention is listed below and no rights to the invention are held by any person, other than the inventor, who would not qualify as an independent inventor under 37 CFR 1.9(c) if that person made the invention, or by any concern which would not qualify as a small business concern under 37 CFR 1.9(d), or a nonprofit organization under 37 CFR 1.9(e). NOTE: Separate verified statements are required from each named person, concern or organization having rights to the invention availing to their status as small entities. (37 CFR 1.27)

Full Name: University of New Orleans Foundation

Address: Office of Research, Administration Building 208, New Orleans, LA 70148

☐ INDIVIDUAL ☒ SMALL BUSINESS CONCERN ☐ NONPROFIT ORGANIZATION

I acknowledge the duty to file, in this application or patent, notification of any change in status resulting in loss of entitlement to small entity status when any new rule 53 application is filed or prior to paying, or at the time of paying, the earliest of the issue fee or any maintenance fee due after the date on which status as a small entity is no longer appropriate. (37 CFR 1.28(b))

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application, any patent issuing thereon, or any patent on which this verified statement is directed.

Name: Mary Louise Trammell

Title: Associate Vice Chancellor for Research/ Director, Office of Technology Transfer

Address: University of New Orleans, Administration Building 208, New Orleans, LA 70148

Signature: Mary L. Trammell

Date: 09/08/99

APPLICATION
FOR
UNITED STATES LETTERS PATENT

TITLE: CATALYST COMPLEX WITH CARBENE LIGAND

APPLICANT: Steven P. Nolan and Jinkun Huang

"EXPRESS MAIL" Mailing Label Number E137209918905

Date of Deposit September 9, 1999
I hereby certify under 37 CFR 1.10 that this correspondence is being deposited with the United States Postal Service as "Express Mail Post Office To Addressee" with sufficient postage on the date indicated above and is addressed to the Assistant Commissioner for Patents, Washington, D.C. 20231.

Francisco Robles
Francisco Robles

CATALYST COMPLEX WITH CARBENE LIGAND

Cross Reference To Related Applications

This application claims priority from U.S.
Provisional Application Serial Nos. 60/099,722, filed
5 September 10, 1998, and 60/115,358, filed January 8, 1999.

Statement as to Federally Sponsored Research

This invention was made with Government support
under Grant No. CHE-963611 awarded by the National Science
10 Foundation. The Government has certain rights in this
invention.

Background of the Invention

The invention relates to metal carbene complexes.
15 More particularly, it relates to catalyst systems
comprising metal carbene complexes.

Catalysts previously known in the art are described
in, for example, U.S. Patent No. 5,312,940 to Grubbs et al.
These catalysts include bis(phosphine) complexes which
20 involve the use of costly phosphine (PR_3) ligands. The
stabilities of such systems, as determined by, for example,
P-C bond degradation at elevated temperature, are limited.
Also, the rates at which bis(phosphine) catalysts carry out
particular reactions are limited. Thus, industrial
25 applications involving large-scale syntheses are not as
efficient as they could be.

Previously available catalytic systems are also
limited in their ability to make highly substituted ring-
closing metathesis (RCM) products. Thus, bis(phosphine)
30 catalysts cannot reliably close dienes to make tri-
substituted cyclic alkenes, and they fail to make tetra-
substituted cyclic alkenes in all but a few cases.

Although Schrock catalysts are available to carry out this type of reaction, such systems are quite sensitive.

Thus there exists in the art a need for a generally air- and moisture-sensitive catalyst system able to carry
5 out RCM reactions efficiently and reliably, and also without excessive thermal sensitivity.

Summary of the Invention

The invention provides catalysts including metal
10 carbene complexes which are useful for synthetic chemical reactions. The catalysts include at least one bulky nucleophilic carbene ligated to the metal center. Methods of making such catalysts, and ligands useful for such catalysts are also provided in the present invention.

15 The inventive catalytic complexes are thermally stable, have high reaction rates, and are air- and moisture-stable. The catalysts of the invention are easy to synthesize, have high catalytic activity, and are relatively inexpensive, due to the availability of the
20 nucleophilic carbene ligand. The catalysts are useful in the facilitation of chemical reactions, including applications in the pharmaceutical industry, fine chemical synthesis, and the synthesis of polymers.

Unless otherwise defined, all technical and
25 scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present
30 invention, suitable methods and materials are described below. In case of conflict, the present specification, including definitions, will control. In addition, the materials, methods, and examples are illustrative only and not intended to be limiting.

Other features and advantages of the invention will be apparent from the following detailed description, and from the claims.

5 Brief Description of the Drawings

Fig. 1A is a general structure of a first particular embodiment of a catalytic complex, having a first ligation pattern.

10 Fig. 1B is a general structure of a first particular embodiment of a catalytic complex, having a second ligation pattern.

Fig. 1C is a general structure of a first particular embodiment of a catalytic complex, having a third ligation pattern.

15 Fig. 2A is an example of a nucleophilic carbene ligand which can be utilized in certain embodiments of the present invention.

Fig. 2B is a particular nucleophilic carbene which can be utilized in certain embodiments of the invention.

20 Fig. 2C is a particular nucleophilic carbene which can be utilized in certain embodiments of the invention.

Fig. 3A is a general structure of a second particular embodiment of a catalytic complex, having a first ligation pattern.

25 Fig. 3B is a general structure of a second particular embodiment of a catalytic complex, having a second ligation pattern.

Fig. 3C is a general structure of a second particular embodiment of a catalytic complex, having a

30 third ligation pattern.

Fig. 4 is an ORTEP diagram of the crystal structure of $\text{Cp}^*\text{Ru}(\text{IMes})\text{Cl}$.

Fig. 5 is an ORTEP diagram of the crystal structure of $\text{Cp}^*\text{Ru}(\text{PCy}_3)\text{Cl}$.

Fig. 6 is an ORTEP diagram of the crystal structure of $\text{Cl}_2\text{Ru}(\text{PCy}_3)(\text{IMes})(=\text{CHPh})$.

Detailed Description

5 The invention includes a catalytic complex for the carrying out of chemical reactions. The complex includes a metal atom and various ligands. A particular embodiment of the catalytic complex is depicted in Figs. 1A, 1B and 1C.

10 Making reference to Fig. 1A, metal atom M can be a transition metal generally having an electron count of from 14 to 18. Particular metals of this description which have been found useful in the present invention include ruthenium and osmium.

15 Ligated to metal atom M are a number of ligands. At least one of these ligands is a carbene ligand, which is functionally an olefin metathesis active fragment, having a carbon atom C^1 which can be further bonded up to two other groups. The bond from metal atom M to carbon atom C^1 can be formulated as the double bonded $\text{M}=\text{C}^1$, although other
20 canonical forms are evidently involved, as detailed in Cotton and Wilkinson's Advanced Inorganic Chemistry, 5th Edition, John Wiley & Sons, New York (1980), pp 1139-1140.

25 As noted, carbon atom C^1 can further bonded to up to two other groups, R and R^1 , and in this case the olefin metathesis active fragment is referred to as an alkylidene. These R and R^1 groups are independently selected from a large number of atoms and substituents. These include hydrogen, alkyl groups having from 1 to 20 carbon atoms (such as methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, and the like). Also possible as either R
30 or R^1 are alkenyl or alkynyl substituents having from 2 to 20 carbon atoms. The groups R and R^1 can also include alkoxy carbonyl substituents having from 2 to 20 carbons atoms, aryl groups, carboxylate substituents having from 1

to 20 carbon atoms, alkoxy substituents having from 1 to 20 carbon atoms, alkenyloxy or alkynyloxy substituents having from 2 to 20 carbon atoms, as well as aryloxy substituents. Also included are alkylthio, alkylsulfonyl, and
5 alkylsulfinyl substituents with from 1 to 20 carbon atoms. Each of the above classes of R or R¹ substituent can be further optionally substituted with halogen, or with alkyl or alkoxy groups of from 1 to 10 carbon atoms, or aryl groups. Further substitution of R and R¹ can include the
10 functional groups of hydroxyl, thiol, thioether, ketone, aldehyde, ester, amide, amine, imine, nitro, carboxylic acid, disulfide, carbonate, isocyanate, carbodiimide, carboalkoxy, carbamate, and halogen.

Any of the above R or R¹ substituents can include
15 various structural isomers (n-, iso-, sec-, and tert-), cyclic or polycyclic isomers, and multiply unsaturated variants.

Particularly useful R and R¹ substituents are vinyl, phenyl, hydrogen, wherein the vinyl and phenyl substituents
20 are optionally substituted with one or more moieties selected from C₁-C₅ alkyl, C₁-C₅ alkoxy, phenyl or a functional group, such as chloride, bromide, iodide, fluoride, nitro, or dimethylamine.

When carbon atom C¹ is not directly bonded to two
25 groups R and R¹, it is further bonded to another carbon C², which is in turn bonded to previously described substituents R and R¹, and the olefin metathesis active carbene ligand is referred to as a vinylidene. This is shown in Fig. 1B. This ligation is generally achieved by
30 means of a double bond from C¹ to C².

Also, as shown in Fig. 1C, C² can be further bonded to another carbon C³. This type of olefin metathesis active carbene ligand is referred to as an allenylidene. C³ is further bonded to the above-described substituents R and R¹.

Carbons C^1 , C^2 and C^3 are each sp^2 hybridized carbons, and the absence of one or two of such carbons in the allenylidene structure of Fig. 1C gives the respective vinylidene or alkylidene or Fig. 1B or 1A, respectively.

5 It has been found that when R or R^1 are aryl, the allenylidene ligand can undergo a rearrangement, forming a different structure in which a ring is formed between C^1 and an aryl carbon of R or R^1 . For example, if $C^1=C^2=C^3Ph_2$ is ligated to metal M in the systems described herein, the
10 olefin metathesis active carbene ligand is not an allenylidene, but rather a cyclized vinyl carbene, an "indenylidene" (in this case phenylindenylidene).

Also ligated to metal atom M are ligands X and X^1 which are anionic ligands, shown in Figs. 1A, 1B and 1C.
15 Such anionic ligands include those independently chosen from halogen, benzoate, C_1 - C_5 carboxylate, C_1 - C_5 alkoxy, phenoxy, and C_1 - C_5 alkylthio groups. In other particular embodiments, X and X^1 are each halide, CF_3CO_2 , CH_3CO_2 , CFH_2CO_2 , $(CH_3)_3CO$, $(CF_3)_2(CH_3)CO$, $(CF_3)(CH_3)_2CO$, PhO, MeO, EtO,
20 tosylate, mesylate, brosylate, or trifluoromethanesulfonate. In other particular embodiments, both X and X^1 are chloride. Ligands X and X^1 can further be bonded to each other, forming a bidentate anionic ligand. Examples include diacid salts, such as dicarboxylate salts.
25 As discussed herein, such groups can alternatively be further bound to a solid phase, for example a polymer support.

Also ligated to metal atom M are ligands L and L^1 . These ligands are chosen from a number of different
30 chemical classes.

One of these classes of ligands L or L^1 is the class of nucleophilic carbenes. In the inventive catalytic complexes, at least one of the ligands L or L^1 is a member of this class. Nucleophilic carbenes are those molecules

having a carbon atom which bears a lone pair of electrons,
desirably also including those molecules additionally
having electron-withdrawing character manifested in atoms
or substituents in electronic communication with, or bonded
5 to, the carbon bearing the lone pair. Such electron
withdrawing atoms or substituents can include atoms which
are more electronegative than carbon, such as nitrogen,
oxygen, and sulfur. These atoms can either be bonded
directly to the carbene carbon, or in a conjugated or
10 hyperconjugated position with respect to this carbon.
Substituents which have electron-withdrawing character
include nitro, halogen, sulfonate, carbonate, sulfide,
thioether, cyano, and other groups known to those in the
art.

15 In particular embodiments, it has been found to be
desirable that not both of ligands L and L¹ be nucleophilic
carbenes, although embodiments in which both L and L¹ are
nucleophilic carbenes are also operative.

Particularly desirable are nucleophilic carbene
20 ligands further substituted with substituents which
increase the steric crowding around the carbon bearing the
lone pair of electrons. These groups can be bonded
directly to the carbene carbon, within a few atoms of the
carbene carbon, or remotely from the carbene carbon, as
25 long as the bulky group is able to inhibit the approach of
agents which tend to react with, and destroy the carbene,
and consequently disable the catalytic complex as a whole.
Thus the stability of the nucleophilic carbene ligand, and
the catalyst itself are fostered by the presence of bulky
30 groups which are able to shield the nucleophilic carbene
from reaction. It should be noted that the olefin
metathesis active carbene fragment is sterically protected
from bimolecular decomposition by the large steric umbrella
provided by the bulky nucleophilic carbene ligand.

Although the invention is not limited by any particular mechanistic theory, it is believed that such a substituent arrangement can provide steric protection from carbene degradation pathways, including thermally induced degradation. The steric bulk of nucleophilic ligands as described herein can lead to more thermally stable catalysts. Such bulky or sterically hindering groups include branched alkyl groups, aryl groups, and aryl groups having branched alkyl substituents, particularly at the ortho positions of the aryl rings. For example, a nucleophilic carbene ligand having bulky alkyl groups such as tert-butyl, iso-propyl or aryl groups with bulky alkyl groups such as 2,4,6-trialkylphenyl or 2,6-dialkylphenyl interacting with the carbene, could be employed in the present invention. The groups L and L¹ can also be further bonded to each other, forming a bidentate ligand wherein either one or both of L and L¹ are nucleophilic carbene ligands.

Cyclic nucleophilic carbene ligands are also envisioned. These may have heteroatoms either in the ring, or bonded to the ring. Particularly desirable examples of this type of nucleophilic carbene ligand are those ligands having a carbene carbon between heteroatoms. Examples include dinitrogen rings such as imidazole, disulfur rings such as 1,3-dithiolane, and dioxygen rings such as 2H,4H-1,3-dioxine. The aromatic, non-aromatic, saturated or unsaturated analogs can be used as well.

Fig. 2A depicts an example of a nucleophilic carbene ligand which can be utilized in certain embodiments of the present invention. Shown is an imidazol-2-ylidene having substituents Y and Y¹, and Z and Z¹. Each substituent is independently selected from a number of carbon-containing groups, or from hydrogen. The carbon-containing groups which can comprise Y, Y¹, Z and Z¹ include alkyl groups

having from 1 to 20 carbon atoms (such as methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, and the like). Also possible are alkenyl or alkynyl substituents having from 2 to 20 carbon atoms. The groups can also
5 include alkoxy carbonyl substituents having from 2 to 20 carbons atoms, aryl groups, carboxylate substituents having from 1 to 20 carbon atoms, alkoxy substituents having from 1 to 20 carbon atoms, alkenyloxy or alkynyloxy substituents having from 2 to 20 carbon atoms, as well as aryloxy
10 substituents. Each of the above classes of substituent can be further optionally substituted with halogen, or with alkyl or alkoxy groups of from 1 to 5 carbon atoms.

Any of the above substituents can include all structural isomers (n-, iso-, sec-, and tert-), cyclic or
15 polycyclic isomers, and multiply unsaturated variants. It should also be noted that the presence of the double bond in the imidazole ring is not required for catalytic activity in the present invention. In certain embodiments, an imidazolidin-2-ylidene can be used as nucleophilic
20 carbene ligand L or L¹.

The structure in Fig. 2B is a particular example of a useful nucleophilic carbene ligand, having both Y and Y¹ as 2,4,6-trimethylphenyl, and both Z and Z¹ as hydrogen. This particular ligand is referred to as 1,3-bis(2,4,6-trimethylphenyl)imidazol-2-ylidene, or IMes. Another
25 example of a useful nucleophilic carbene is given in Fig. 2C, which shows a structure having both Y and Y¹ as 2,6-diisopropylphenyl, and both Z and Z¹ as hydrogen. This particular ligand is referred to as 1,3-bis(2,6-diisopropylphenyl)imidazol-2-ylidene, or IPr.
30

Another class of ligand which can serve as L or L¹ is the class of phosphines. Particularly useful are trialkyl- or triarylphosphines, such as trimethylphosphine, triphenylphosphine, triisopropylphosphine, and similar

phosphines. The phosphines tricyclohexylphosphine and tricyclopentylphosphine are also useful, and are collectively referred to as PCy_3 .

Other classes of ligands which can serve as L or L^1 are sulfonated phosphines, phosphites, phosphinites, phosphonites, arsine, stibine, imines, ethers, amines, amides, sulfoxides, carbonyls, carboxyls, nitrosyls, pyridines, and thioethers.

Other embodiments of catalytic complexes useful in the present invention are shown in Figs. 3A (alkylidene), 3B (vinylidene) and 3C (allenylidene), in which the analogy with the series of Figs. 1A, 1B and 1C is based on the identity of the olefin metathesis active carbene ligand, alkylidene, vinylidene and allenylidene, respectively. The elements M, X, C^1 , C^2 , C^3 , R and R^1 are as described above for the first described embodiment of the inventive catalytic complex. In this second particular embodiment, ligand L is a nucleophilic carbene ligand, as described above. In addition, since the species depicted in Figs. 3A, 3B, and 3C are all cationic complexes, an anion A^- is required. This anion can be any inorganic anion, and can also include some organic anions. Thus, A^- can be, for example, halide ion, SbF_6^- , PF_6^- , BF_4^- , AsCl_4^- , O_3SONO^- , SO_2F^- , NSO_3^- , azide, nitrite, nitrate, or acetate, and many others known to those of skill in the art.

In this embodiment, another ligand of metal M is Ar, which is an aromatic ring system, including the η^6 -bonded system. The symbol η is used to signify that all aromatic ring atoms are bonded to the metal atom. Such systems include C_6H_6 ring systems, and various alkyl substituted C_6H_6 ring systems. Heterocyclic arene rings are also suitable, and these include $\eta^6\text{-C}_5\text{H}_5\text{N}$, and alkyl substituted derivatives thereof. These rings can have substituents chosen from a wide range of groups including alkyl groups

having from 1 to 20 carbon atoms (such as methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, and the like). Also possible are alkenyl or alkynyl substituents having from 2 to 20 carbon atoms. The groups can also
5 include alkoxy carbonyl substituents having from 2 to 20 carbons atoms, aryl groups, carboxylate substituents having from 1 to 20 carbon atoms, alkoxy substituents having from 1 to 20 carbon atoms, alkenyloxy or alkynyloxy substituents having from 2 to 20 carbon atoms, as well as aryloxy
10 substituents. Each of the above classes of substituent can be further optionally substituted with halogen, or with alkyl or alkoxy groups of from 1 to 5 carbon atoms. For example, useful η^6 -bonded L or L¹ ligands are p-cymene, fluorene and indene.

15 The inventive catalytic complexes can be used as homogeneous catalysts, or are equally well suited as heterogeneous catalysts. The latter embodiment is realized by linking the catalytic complexes to a suitable solid phase, such as a polymeric support. The solid phase can be
20 bound to the catalytic complex either cleavably or non-cleavably. The solid phase can be a polymer which can either be a solid-state resin such as a Wang resin, or a soluble polymer such as non-crosslinked chloromethylated polystyrene (NCPS). This polymer shows excellent
25 properties, such as solubility in tetrahydrofuran (THF), dichloromethane, chloroform, and ethyl acetate, even at low temperatures (-78°C). NCPS is insoluble in water and methanol. These features allow traditional organic chemistry techniques such as solvent extraction, and
30 methanol precipitation. Suitable polymers include hydroxyl-containing polymers such as Wang resin, or poly(ethylene glycol) (PEG).

The method of attachment between solid phase and catalytic complex can take the form of a link to the ligand

L or L¹, which is desirably the nucleophilic carbene ligand. This arrangement is desirable since the catalytic complex is believed to operate by first releasing the ligand which is not a nucleophilic carbene, for example, by releasing a phosphine ligand. Thus, linkage to the phosphine ligand would result in loss of the solid phase-catalytic complex interaction, upon catalysis. Also considered desirable is linkage of the catalytic complex to a solid phase through the anionic ligands X and/or X¹. Thus, any linkage which involves a group serving as an anionic ligand as described above can be used to attach the catalytic complex to a solid support. For example, carboxylate resins can be employed for this purpose.

The inventive catalytic complexes are air- and moisture-stable, and thus can be used under atmospheric conditions, and even in aqueous environments. The stability of the catalytic substrates and products will be the limiting factors with respect to use under such conditions. The inventive catalytic complexes are soluble in typical organic solvents, such as tetrahydrofuran (THF), benzene, toluene, xylene, diethyl ether, dioxane, alcohols, acetonitrile, dimethylsulfoxide (DMSO), dimethylformamide (DMF), and similar solvents, but not particularly soluble in water or methanol.


The catalytic complexes need not be used in the presence of any initiators or cocatalysts, although materials such as phosphine sponges can optionally be used. Those of skill in the art will recognize the identity of the members of this class, which includes copper chloride, and Lewis acids generally, in concentrations up to those stoichiometric with that of the catalytic complex.

Use of Catalytic Complexes in Ring-Closing Metathesis (RCM)

The catalytic complexes can be used for ring-closing metathesis. This reaction converts a diterminal diene (a compound having two $-C^a=C^bH_2$ groups, the C^a atoms of which are able to link together to form a cyclic compound with a $-C^a=C^a-$ linkage), to a cyclic alkene, with $H_2C^b=C^bH_2$ as a side product. In some instances, the diterminal diene (or an α, ω diene) can undergo a 1,3-hydrogen shift rearrangement (to give an $\alpha, \omega-1$ diene), and the product will be a cyclic alkene with one less methylene group in the ring, and propene as a side product.

A pronounced solvent dependence of the reactivity of the present catalytic complexes was noticed. As can be seen from the results compiled in Scheme 1, reaction rates for $(IMes)(Pcy_3)Cl_2Ru(=CHPh)$ in toluene are substantially higher than those in CH_2Cl_2 (the substituent E is $-CO_2Et$). Thus, the tetrasubstituted cyclohexene derivative of Scheme 1 is formed in essentially quantitative yield after only 15 min if the reaction is carried out in toluene. The reaction requires 2-3 hours in CH_2Cl_2 to reach completion. This influence of the reaction medium has been observed for the ruthenium carbene complexes bearing *N-mesityl* substituents on their imidazol-2-ylidene ligands. However, the related complexes having *N-cyclohexyl* or *N-isopropyl* groups do not show this effect.


Scheme 1. Solvent Dependence of the Reactivity of Complex
 (IMes) (Pcy₃) Cl₂ Ru (=CHPh)



This reactivity of (IMes) (Pcy₃) Cl₂ Ru (=CHPh) in toluene is impaired by a tendency of the active species to promote isomerization of the double bonds of the substrate. Thus, in Scheme 2, treatment of the pictured diene with as little as 1.2 mol% of (IMes) (Pcy₃) Cl₂ Ru (=CHPh) in toluene leads to complete consumption of the starting material within 45 min, but delivers significant amounts of the 20-membered ring in addition to the desired 21-membered lactone.














Although not wishing to be bound by any particular theory, the cis-cyclic alkene is believed to result from an initial isomerization of one of the double bonds in the starting material, followed by elimination of propene instead of ethylene during ring closure. This intrinsic bias for ring contraction was not suppressed by lowering the reaction temperature. In stark contrast, however, only minute amounts of the cis-alkene are detected if the reaction is performed in CH₂Cl₂.

Scheme 2. Stereochemistry



As can be seen from the results compiled in Table 1, the reactivities of (IMes)(Pcy₃)Cl₂Ru(=CHPh) and (IMes)(Pcy₃)Cl₂Ru(phenylindenylidene) in CH₂Cl₂ are sufficiently high to allow the preparation of di-, tri- and even tetrasubstituted cyclo-alkenes in good to excellent yields. All ring sizes including medium and macrocyclic ones can be accessed. The yield data given are the isolated yields. The reactions with yields given with superscript b (entries 1-4) were carried out in toluene at 80°C. The compound 3a refers to (IMes)(Pcy₃)Cl₂Ru(=CHPh), and 3b to (IMes)(Pcy₃)Cl₂Ru(phenylindenylidene). E is -CO₂Et.

Table 1. RCM catalyzed by (IMes)(Pcy₃)Cl₂Ru(=CHPh) and (IMes)(Pcy₃)Cl₂Ru(phenylindenylidene) in CH₂Cl₂.

Entry	Product	Catalyst (mol%)	Yield (%)
1		3a (2%)	96 ^b
2		3b (2%)	97 ^b
3	 (E=CO ₂ Et)	3a (5%)	77 ^b
4		3b (2%)	89 ^b
5		3a (5%)	98
6		3a (5%)	93
7		3b (5%)	71
8		3a (1%)	64
9		3a (1%)	62 (R = H)
10		3a (5%)	95 (R = Me)
11		3a (2%)	72
12		3a (3%)	82
13		3a (4%)	71

5 It must be noted that most of these cyclizations cannot be carried out if the bis(phosphine) complex

(PCy₃)₂Cl₂Ru(=CHPh) is used as the catalyst. This holds true for all tetrasubstituted cases (entries 1-4 and 7), the trisubstituted 8-membered ring shown in entry 10, as well as for annulation reactions depicted in entries 5 and 6. Although the macrocyclic products (entries 11-13) can also be obtained with the use of (PCy₃)₂Cl₂Ru(=CHPh), using (IMes)(PCy₃)Cl₂Ru(=CHPh) results in shorter reaction times and allows lower catalyst loadings to be employed. This aspect is particularly relevant with respect to pentadec-10-enolide (entry 11) which is converted into the valuable, musk-odored perfume ingredient EXALTOLIDE[®] (= pentadecanolide) upon simple hydrogenation.

As can be deduced from the results in Table 1, complex (IMes)(PCy₃)Cl₂Ru(=CHPh) bearing a benzyldiene carbene moiety and complex (IMes)(PCy₃)Cl₂Ru(phenylindenylidene) with a phenylindenylidene unit are essentially equipotent pre-catalysts.

Method of Making Catalytic Complexes

The inventive catalytic complexes can be made according to the following general synthetic procedures, which are adapted from known procedures.

To synthesize a catalytic complex according to a first embodiment of the invention, one of the two phosphine ligands of a diphosphine-ligated ruthenium or osmium catalyst is exchanged with a nucleophilic carbene ligand. For example, starting material diphosphine-ligated complexes (PCy₃)Cl₂Ru(=CHPh) and (PPh₃)Cl₂Ru(=CHPh) can be synthesized according to general procedures such as those given by Schwab et al., *Angew. Chem. Intl. Ed. Engl.*, (1995) 34, 2039-41.

Ligand-exchange reactions are carried out by exposing the diphosphine-ligated complexes to nucleophilic carbene ligands, as defined above, in suitable solvents such as THF, toluene, and the like. Reactions are
5 generally carried out at temperatures of from about 0°C to about 50°C, for about 15 minutes to several hours. Subsequent recrystallization in inert solvents gives the complexes in good yield and high purity.

The nucleophilic carbene ligands according to the
10 invention are synthesized according to the following general synthetic procedure. Solutions of heteroatom-containing starting material such as aniline, or substituted aniline, phenol or substituted phenol, benzenethiol or substituted benzenethiol, primary- or
15 secondary-amines, alcohols and thiols can be prepared in solvents such as tetrahydrofuran (THF), benzene, toluene, xylene, diethyl ether, dioxane, alcohols, acetonitrile, dimethylsulfoxide (DMSO), dimethylformamide (DMF), water, and similar solvents, under an inert atmosphere.
20 Substituents for the above groups include alkyl groups having from 1 to 20 carbon atoms (such as methyl, ethyl, n-propyl, iso-propyl, n-butyl, iso-butyl, sec-butyl, and the like). Also possible are alkenyl or alkynyl substituents having from 2 to 20 carbon atoms. The groups can also
25 include alkoxycarbonyl substituents having from 2 to 20 carbons atoms, aryl groups, carboxylate substituents having from 1 to 20 carbon atoms, alkoxy substituents having from 1 to 20 carbon atoms, alkenyloxy or alkynyloxy substituents having from 2 to 20 carbon atoms, as well as aryloxy
30 substituents. Each of the above classes of substituent can be further optionally substituted with halogen, or with alkyl or alkoxy groups of from 1 to 5 carbon atoms. Particularly useful are those substituents such as methyl, ethyl, propyl, and butyl, including branched isomers, and

aryl substituents at the *ortho*- or *diortho*-positions (for example, 2- or 2,6- substitution for benzyl rings).

The solution is then contacted with an approximately one half of equimolar amount (with respect to the heteroatom-containing starting material) of paraformaldehyde. After heating to dissolve paraformaldehyde, the contents of the flask are acidified with an approximately one half of equimolar amount (with respect to the heteroatom-containing starting material) of mineral acid (for example, hydrochloric acid or nitric acid).


At this stage, if a nitrogen-containing starting material (aniline-derivative or primary amine-derivative) is used, an approximately one half of equimolar amount (with respect to the heteroatoms-containing starting material) of a dialkoxyacetaldehyde is added drop wise after a few minutes of stirring. The dialkoxyacetaldehyde can be dimethoxy-, diethoxy-, dipropoxy-, dibutoxy-, diphenoxy, or can be any of a number of combinations of such alkoxy substituents such as for example methoxyethoxy, or methoxyphenoxy. The procedure then continues as follows.

If, on the other hand, oxygen or sulfur heteroatom-containing starting material is used, the above paragraph is not followed, and the procedures from this point on are common to all starting materials. After equipping the reaction flask with a Dean-Stark trap, or similar device, the mixture is heated to a temperature of from about 80°C to about 180°C, preferably from about 100°C to about 150°C for several hours (from about 5 to about 30 hours). During this time, a precipitate forms, as the side products of water and methanol, as well as some solvent, are removed. The reaction mixture is stirred at room temperature for a time ranging from about 20 minutes to about 4 hours,

preferably from 1 to 3 hours. Precipitate will have formed during this time.

The precipitate is filtered, washed with a suitable solvent, such as THF to give the nucleophilic carbene product in the form of a salt. For example, if aniline or substituted aniline is used, the product will be a 1,3-diarylimidazole salt. If the starting material is a primary amine, the product will be a 1,3-dialkyl imidazole salt. Either of these products can be converted to the saturated heterocyclic derivative (imidazolidine) by conventional hydrogenation techniques such as exposure to H₂ over a carbon-palladium or carbon-platinum catalyst. Such techniques will be recognized and known to those of skill in the art. If the starting material is a phenol- or thiobenzene-derived compound, the product will be a dibenzoxymethane-, or dibenzthiomethane-product. If the starting material is an alcohol or thiol, the product will be a 1,1-bis(alkoxy)methane- or 1,1-bis(alkylthio)methane-product.

The second embodiments of the catalytic complexes of the invention are easily made by combining a precursor species of the catalytic complexes with an acetylene to give the allenylidene type of catalytic complex (see Fig. 3C). An example of this precursor species of the catalytic complex is shown below.



In this structure, metal M, and ligands X, X¹, L and Ar are defined as above, with L being a nucleophilic carbene. This precursor species is generally available in

the form of a dimer $[\text{ArRuCl}_2]_2$, which is converted to the precursor species when the dimer is exposed to a nucleophilic carbene in a suitable solvent such as THF, hexanes and other non-protic solvents. For example, the
5 dimer $[(p\text{-cymene})\text{-RuCl}_2]_2$ is commercially available from Strem Chemicals (Newburyport, MA).

The acetylenes with which precursor species of the inventive catalytic complexes combine to form second
embodiments of the invention are terminal acetylenes, and
10 can be substituted at the γ -position with alkyl or aryl groups, or optionally further substituted with halogen, or with alkyl or alkoxy groups of from 1 to 10 carbon atoms, or aryl groups. Further substitution can include the functional groups of hydroxyl, thiol, thioether, ketone,
15 aldehyde, ester, amide, amine, imine, nitro, carboxylic acid, disulfide, carbonate, isocyanate, carbodiimide, carboalkoxy, carbamate, and halogen. Particularly useful substituents are vinyl, phenyl, or hydrogen, wherein the vinyl and phenyl substituents are optionally substituted
20 with one or more moieties selected from $\text{C}_1\text{-C}_5$ alkyl, $\text{C}_1\text{-C}_5$ alkoxy, phenyl or a functional group, such as chloride, bromide, iodide, fluoride, nitro, or dimethylamine.

The invention will be further described in the following examples, which do not limit the scope of the
25 invention described in the claims.

Examples

Illustrations of methods of making certain
embodiments of the inventive catalytic complexes, as well
30 as properties thereof, are provided by the following examples.

Example 1: Synthesis of IMes-HCl

A 300 mL Schlenk flask was charged with 2,4,6-trimethylaniline (10g, 74 mmol), toluene (50 mL), and paraformaldehyde (1.11 g, 37 mmol) under argon and heated to 110°C until all the paraformaldehyde was dissolved. The flask was then cooled to 40°C and HCl (6.17 mL, 6N, 37 mmol) was added to the reaction mixture drop wise. The mixture was stirred at that temperature for 10 minutes before dimethoxyacetaldehyde (6.442 g, 60% wt. in water, 37 mmol) was added in drop wise fashion. The flask was then equipped with a Dean-Stark trap and heated to 120°C for 15 hours, during which time a dark precipitate was formed and grew in volume by removal of the side-products (H₂O and methanol) and some of the solvent through the Dean-Stark trap. The reaction mixture was then allowed to cool to room temperature and stirred at that temperature for two hours. Filtration of the precipitate through a Schlenk frit, washing with tetrahydrofuran (three times, 20 mL each wash), and drying yielded a white solid in 60% yield, which was characterized spectroscopically as pure IMes-HCl. ¹H NMR: δ = 2.12 (s, 12H, o-CH₃), 2.30 (s, 6H, p-CH₃), 6.97 (s, 4H, mesityl), 7.67 (s, 2H, NCHCHN), 10.68 (s, 1H, HCl).

Example 2: Synthesis of IMes

In a glovebox, a 300 mL Schlenk flask equipped with a stir bar was charged with 20.0g (58.7 mmol) of IMes-HCl and 120 mL of dry tetrahydrofuran. The resulting suspension was stirred for 10 minutes after which time 6.80 g (60.7 mmol) of solid potassium tert-butoxide was added to the suspension at room temperature in a single portion. A dark gray solution was obtained immediately. The flask was taken out of the glovebox and connected to the Schlenk line. The solution was stirred for 20 minutes before all

volatiles were removed under vacuum. The residue was extracted into warm toluene (120 mL + 60 mL + 20 mL) and filtered through a medium porosity frit (filtration was rather slow), and the solvent was removed under vacuum to obtain crystals of IMes. The resulting product was recovered in 90% yield, and had a dark tint but was sufficiently pure for its use in further synthesis. Further purification could be achieved by recrystallization from toluene or hexane, yielding colorless crystals.

The synthesis of related carbenes 1,3-bis(4-methylphenyl)imidazol-2-ylidene (ITol) and 1,3-bis(4-chlorophenyl)imidazol-2-ylidene (IpCl) was carried in an analogous fashion.

Example 3: Synthesis of (IMes)(PCy₃)(Cl)₂Ru(=CHCH=CMe₂)

The procedure was carried out under purified and dried argon atmosphere and with dried and degassed solvents. IMes (2.1990 g, 7.221 mmol) was suspended in 250 mL hexanes, into which (Cl)₂(PCy₃)₂Ru(=CHCH=CMe₂) (5.0718 g, 7.092 mmol) was added in one portion. The mixture was heated for 2.5 hours with stirring at 60°C. During this period, the formation of an orange-brown precipitate was observed. The volume of the suspension was then reduced in vacuum to 50 mL and the suspension was cooled to -78°C.

Following filtration and cold pentane washing of the residue (2 washes, each 20 mL), the product was isolated as a brown orange microcrystalline material in 72% yield (3.97g).

Example 4: Synthesis of (IMes)(PCy₃)Cl₂Ru(=CHPh)


The procedure of Example 3 was followed, except that (Cl)₂(PCy₃)₂Ru(=CHPh) was used. This complex was soluble in a variety of organic solvents including hydrocarbon, tetrahydrofuran, acetone, methylene chloride, and

diethylether. The identity of the complex was confirmed by X-ray crystallography. Other embodiments will be readily synthesized by substituting the IMes ligand with other nucleophilic carbene ligands.

5

Example 5: Thermodynamic Studies

The thermodynamics of the following reaction in tetrahydrofuran (THF) at room temperature were studied.



(Cp* is $\eta^5\text{-C}_5\text{Me}_5$) The reaction proceeds rapidly as indicated by the rapid development of a deep blue color in the reaction solution. A deep blue crystalline solid was
15 isolated in 86% yield. Nuclear magnetic resonance data of the blue solid indicated the isolation of a single species bearing a unique Cp* and a single carbene ligand. X-ray crystallography confirmed the formulation of Cp*Ru(IMes)Cl. An enthalpy of reaction of -62.6 ± 0.2 kcal/mol was
20 measured by anaerobic solution calorimetry in THF at 30°C when 4 equivalents of carbene were reacted with 1 equivalent of the tetramer, [Cp*RuCl]₄. Table 1 compares the enthalpy of similar reactions where IMes is replaced with other moieties.


25

Table 1. Comparison of Reaction Enthalpies

Identity of L in $\text{Cp}^*\text{Ru}(\text{L})(\text{Cl})$	ΔH (kcal/mol) of reaction: $[\text{Cp}^*\text{RuCl}]_4 + 4 \text{ L} \rightarrow 4$ $\text{Cp}^*\text{Ru}(\text{L})(\text{Cl})$	Relative stability of Ru-L bond (kcal/mol)
Imes	-62.6 ± 0.2	-15.6
$\text{P}(\text{isopropyl})_3$	-37.4 ± 0.3	-9.4
$\text{P}(\text{cyclohexyl})_3$	-41.9 ± 0.2	-10.5

The IMes ligand proves to be a stronger binder to the Cp^*RuCl fragment than PCy_3 , by 5 kcal/mol. The carbene ligand is a fairly good binder but can be displaced if a better donor ligand, such as a phosphite, is used. The phosphite reaction allows for the construction of a thermochemical cycle which confirms the internal consistency of the calorimetric data, as shown in Scheme 1.

Scheme 1. Thermodynamic Cycle



A further verification of the thermochemical results can be made by examining the following hypothetical reaction.



This reaction is calculated to be exothermic by 5 kcal/mol and no entropic barrier is apparent, so the reaction should proceed readily as written. Indeed, upon
 10 mixing of the reagents in THF- d_8 , the characteristic ^{31}P signal of $\text{Cp}^*\text{Ru}(\text{PCy}_3)\text{Cl}$ disappears (at 11.3 ppm), and that of free PCy_3 appears (40.4 ppm), as observed by Campion et al., *J. Chem. Soc. Chem. Commun.*, (1988) 278-280.

15 Example 6: Structural Studies

In order to gauge the steric factor inherent in the catalytic systems, structural studies were carried out on $\text{Cp}^*\text{Ru}(\text{IMes})\text{Cl}$ (Fig. 4), $\text{Cp}^*\text{Ru}(\text{PCy}_3)\text{Cl}$ (Fig. 5), and $(\text{IMes})(\text{PCy}_3)\text{Cl}_2\text{Ru}(=\text{CHPh})$ (Fig. 6). Comparison was made to
 20 another sterically demanding ligand in the complex $\text{Cp}^*\text{Ru}(\text{P}^i\text{Pr}_3)\text{Cl}$. The following crystal data was obtained. For $\text{Cp}^*\text{Ru}(\text{IMes})\text{Cl}$: monoclinic, space group $\text{P2}_1/\text{c}$, dark blue prism, $0.35 \times 0.25 \times 0.20$, $a = 10.6715$ (2), $b = 14.3501$ (3), $c = 19.2313$ (4), $\beta = 103.2670$ (10) deg, $Z = 4$, $R_f = 0.0294$, $\text{GOF} =$
 25 0.888 . For $\text{Cp}^*\text{Ru}(\text{PCy}_3)\text{Cl}$: orthorhombic, space group Pcba , dark blue prism, $0.45 \times 0.35 \times 0.25$, $a = 18.9915$ (6), $b = 15.6835$ (5), $c = 19.0354$ (6), $Z = 8$, $R_f = 0.0392$, $\text{GOF} = 1.132$. For $(\text{IMes})(\text{PCy}_3)\text{Cl}_2\text{Ru}(=\text{CHPh})$: space group $\text{P2}_12_12_1$, yellow-orange prism, $a = 12.718$ (1), $b = 14.549$ (1), $c = 26.392$ (2),
 30 $R_f = 0.0616$, $z = 4$, $\text{GOF} = 1.038$. The metrical data of $\text{Cp}^*\text{Ru}(\text{P}^i\text{Pr}_3)\text{Cl}$ (Campion et al., *J. Chem. Soc. Chem. Commun.*, (1988) 278-280) can be used for comparison: Ru-P , 2.383 (1) Å; Ru-Cl , 2.378 (1) Å; $\text{Ru-Cp}^*(\text{c})$, 1.771 (1) Å; Cl-Ru-P ,

91.2 (1)°; Cl-Ru-Cp*(c), 129.9 (1)°; C(1)-Ru-Cp*(c), 139.9 (1)°.

The three Cp*RuCl(L) structures are similar, with the variation in Ru-L distances the only standout feature, but this is explainable by the difference in covalent radii between P and C. Only slight angle distortions are observed in Cp*Ru(IMes)Cl, presumably to accommodate the bulkiness of IMes. The IMes ligand displays non-coplanar rings with torsion angles of 78.46 (4)° between the arene ring bound to N(2) and the imidazole ring and 78.78 (5)° between the imidazole ring and the arene ring bound to N(1). The two arene rings adopt a mutually staggered configuration.

A direct comparison of the steric properties displayed by IMes and PCy₃ provides insight into the significant steric congestion provided by the IMes ligation. The cone angle reported for PⁱPr₃ and PCy₃ are 160° and 170°, respectively (Tolman, *Chem.Rev.* (1977) 77, 313-348). Such a cone angle measurement is not straightforward in the present system. Instead, the crystallographic data can be used to determine closest contact angles involving non-hydrogen atoms in Cp*Ru(IMes)Cl and Cp*Ru(PCy₃)Cl. For the Ru-PCy₃ fragment, an angle of 96.3° is measured using cyclohexyl methylene carbons on adjacent cyclohexyl rings defining the largest angle. For the Ru-PⁱPr₃ fragment in Cp*Ru(PⁱPr₃)Cl a similar angle of 95.8° is obtained. As for the IMes fragment, two parameters can be obtained. Angles of 150.7° and 115.3° are measured for the <4-Me-Ru-4'-Me and <6-Me-Ru-2'-Me angles, respectively. The steric coverage of the IMes ligand can be considered as a fence rather than a cone. The increased steric congestion provided by the IMes ligand compared to PCy₃ derives from the presence of bulky substituents on the imidazole nitrogens and, to a greater extent, from the significantly shorter metal-carbon bond

distance which brings the entire IMes ligand closer to the metal center.

The structural analysis of (IMes)(PCy₃)Cl₂Ru(=CHPh) shown in Fig. 6 reveals a distorted square pyramidal

5 coordination with a nearly linear Cl(1)-Ru-Cl(2) angle (168.62°). The carbene unit is perpendicular to the C(1)-Ru-P plane, and the carbene aryl moiety is only slightly twisted out of the Cl(1)-Ru-Cl(2)-C(40) plane. The Ru-C(40) bond distance (1.841 (11)Å) is the same as that in
10 RuCl₂(=CH-*p*-C₆H₄Cl)(PCy₃)₂ (1.838 (3)Å) and shorter than that in (PCy₃)₂RuCl₂(=CHCH=CPh₂) (1.851 (21)Å). While two (formally) carbene fragments are present in (IMes)(PCy₃)Cl₂Ru(=CHPh), they display different Ru-C distances (Ru-C(40) = 1.841 (11) and Ru-C(1) = 2.069 (11)Å).
15 These important metrical parameters clearly distinguish two metal-carbene interactions: a metal benzylidene fragment with a formal metal to carbon double bond and a metal imidazolium carbene with a formal metal-carbon single bond. From Fig. 6, it is also clear that the IMes ligand is
20 sterically more demanding than PCy₃.

Example 7: Thermal Stability Studies

In the course of catalytic testing, the remarkable air stability of the inventive catalytic complexes was
25 observed. To gauge the robust nature of these carbene complexes in solution, their thermal stability under inert atmosphere was tested at 60°C. The relative order of stability found was (IMes)(PCy₃)Cl₂Ru(=CHPh) >> (IMes)(PPh₃)Cl₂Ru(=CHPh) > (PCy₃)₂Cl₂Ru(=CHPh). After 14
30 days of continuous heating of toluene solutions of (IMes)(PCy₃)Cl₂Ru(=CHPh) to 60°C, no decomposition was detected (as monitored by both ¹H and ³¹P NMR). In contrast, solutions of (PCy₃)₂Cl₂Ru(=CHPh) showed signs of decompositions after one hour, under the same conditions.

The catalyst (IMes)(PCy₃)Cl₂Ru(=CHPh) was stable at 100°C for 36 hours before showing any indication of decomposition. Similar thermal decomposition studies have been conducted in refluxing methylene chloride,
5 dichloromethane, toluene, benzene and diglyme with similar results.

Other Embodiments

It is to be understood that while the invention has
10 been described in conjunction with the detailed description thereof, the foregoing description is intended to illustrate and not limit the scope of the invention, which is defined by the scope of the appended claims. Other aspects, advantages, and modifications are within the scope
15 of the following claims.

What is claimed is:

1 1. A catalytic complex comprising:

2 a) a metal atom selected from the group

3 consisting of ruthenium or osmium;

4 b) at least one anionic ligand ligated to said
5 metal;

6 c) at least one nucleophilic carbene ligand ligated
7 to said metal;

8 d) a further ligand ligated to said metal; and

9 e) a carbon-containing ligand ligated to said metal,

10 wherein said carbon-containing ligand is

11 selected from the group consisting of

12 alkylidene, benzylidene, indenylidene, vinylidene,

13 and allenylidene.

1 2. The catalytic complex of claim 1, wherein the anionic

2 ligand is selected from the group consisting of halide,

3 carboxylate, alkoxy, aryloxy, and alkyl sulfonate.

1 3. The catalytic complex of claim 2, wherein the anionic

2 ligand is chloride.

1 4. The catalytic complex of claim 1, wherein the

2 nucleophilic carbene ligand comprises a carbene carbon

3 further bonded to two heteroatoms having electronegativity

4 greater than that of carbon, wherein the heteroatoms are

5 independently selected from the group consisting of

6 nitrogen, oxygen, and sulfur.

1 5. The catalytic complex of claim 4, wherein the

2 nucleophilic carbene ligand comprises a substituted or

3 unsubstituted, saturated or unsaturated 1,3-diheteroatomic


4 cyclic compound.

1 6. The catalytic complex of claim 1, wherein the complex
2 is linked to a solid support by means of a link between
3 said anionic ligand and said solid support.

1 7. The catalytic complex of claim 1, wherein the complex
2 is linked to a solid support by means of a link between
3 said nucleophilic carbene and said solid support.

1 8. The catalytic complex of claim 1, wherein the carbon-
2 containing ligand selected from the group consisting of
3 alkylidene, vinylidene and allenylidene is further
4 substituted with substituents selected from the group
5 consisting of hydrogen, C₁-C₂₀ alkyl, C₂-C₂₀ alkenyl, C₂-C₂₀
6 alkynyl, C₂-C₂₀ alkoxy carbonyl, aryl, C₁-C₂₀ carboxylate, C₁-
7 C₂₀ alkoxy, C₂-C₂₀ alkenyloxy, C₂-C₂₀ alkynyloxy, and aryloxy,
8 optionally being substituted with C₁-C₅ alkyl, halogen, C₁-C₆
9 alkoxy, or with a phenyl group substituted with halogen, C₁-
10 C₅ alkyl or C₁-C₅ alkoxy.

1 9. A catalytic complex of the formula:



2
3 wherein M is Os or Ru;
4 C¹, C² and C³ are sp²-hybridized carbons, wherein
5 either or both of C¹ and C² are optionally absent;
6 R and R¹ are independently selected from the group
7 consisting of hydrogen, C₁-C₂₀ alkyl, C₂-C₂₀ alkenyl, C₂-C₂₀
8 alkynyl, C₂-C₂₀ alkoxy carbonyl, aryl, C₁-C₂₀ carboxylate, C₁-
9 C₂₀ alkoxy, C₂-C₂₀ alkenyloxy, C₂-C₂₀ alkynyloxy, or aryloxy,
10 each R and R¹ optionally being substituted with C₁-C₅ alkyl,
11 halogen, C₁-C₆ alkoxy, or with a phenyl group substituted
12 with halogen, C₁-C₅ alkyl or C₁-C₅ alkoxy;

13 X and X¹ are independently selected from the group
14 consisting of anionic ligands; and


15 L and L¹ are selected from the group consisting of
16 nucleophilic carbenes, phosphine, sulfonated phosphine,
17 phosphite, phosphinite, phosphonite, ether, amine, amide,
18 sulfoxide, carbonyl, nitrosyl, pyridine and thioether,
19 wherein at least one of L or L¹ is a nucleophilic carbene.

1 10. The catalytic complex according to claim 9, wherein
2 only one of L or L¹ is a nucleophilic carbene.

1 11. A catalytic complex according to claim 10, wherein one
2 of L or L¹ is a phosphine.

1 12. A catalytic complex according to claim 9, wherein at
2 least 2 of X, X¹, L or L¹ are bonded together to form a
3 multidentate ligand.

1 13. A catalytic complex according to claim 9, wherein said
2 nucleophilic carbene is of the formula:




3
4 wherein

5 Y and Y¹ are independently selected from the group
6 consisting of hydrogen, C₁-C₂₀ alkyl, C₂-C₂₀ alkenyl, C₂-C₂₀
7 alkynyl, C₂-C₂₀ alkoxy carbonyl, aryl, C₁-C₂₀ carboxylate, C₁-
8 C₂₀ alkoxy, C₂-C₂₀ alkenyloxy, C₂-C₂₀ alkynyloxy, or aryloxy,
9 each Y and Y¹ optionally being substituted with C₁-C₅ alkyl,
10 halogen, C₁-C₆ alkoxy, or with a phenyl group substituted
11 with halogen, C₁-C₅ alkyl or C₁-C₅ alkoxy and;

12 Z and Z¹ are independently selected from the group
13 consisting of hydrogen, C₁-C₂₀ alkyl, C₂-C₂₀ alkenyl, C₂-C₂₀

14 alkynyl, C₂-C₂₀ alkoxy carbonyl, aryl, C₁-C₂₀ carboxylate, C₁-
 15 C₂₀ alkoxy, C₂-C₂₀ alkenyloxy, C₂-C₂₀ alkynyloxy, or aryloxy,
 16 each Z and Z¹ optionally being substituted with C₁-C₅ alkyl,
 17 halogen, C₁-C₆ alkoxy, or with a phenyl group substituted
 18 with halogen, C₁-C₅ alkyl or C₁-C₅ alkoxy,
 19 and wherein the ring can be optionally aromatic by
 20 introduction of a further double bond in the ring.

1 14. A catalytic complex of the formula:



2
 3 wherein

4 C¹, C² and C³ are sp²-hybridized carbons, wherein
 5 either or both of C¹ and C² are optionally absent;

6 M is selected from the group consisting of Os and
 7 Ru;

8 R and R¹ are independently selected from the group
 9 consisting of hydrogen, C₁-C₂₀ alkyl, C₂-C₂₀ alkenyl, C₂-C₂₀
 10 alkynyl, C₂-C₂₀ alkoxy carbonyl, aryl, C₁-C₂₀ carboxylate, C₁-
 11 C₂₀ alkoxy, C₂-C₂₀ alkenyloxy, C₂-C₂₀ alkynyloxy, or aryloxy,
 12 each R and R¹ optionally being substituted with C₁-C₅ alkyl,
 13 halogen, C₁-C₆ alkoxy, or with a phenyl group substituted
 14 with halogen, C₁-C₅ alkyl or C₁-C₅ alkoxy;

15 X is an anionic ligand; and

16 L is a nucleophilic carbene; and

17 Ar is an aryl substituent, bonded to M by an η⁶
 18 bond.

1 15. A catalytic complex according to claim 13, wherein one
 2 of L or L¹ is a phosphine.

7 b) heating said first reaction mixture until said
8 paraformaldehyde dissolves;

9 c) adding an approximately one-half equimolar
10 amount of a dialkoxyacetaldehyde to make a second reaction
11 mixture; and

12 d) heating said second reaction mixture for a time
13 and under conditions sufficient to make a nucleophilic
14 carbene.

1 19. The method of claim 18, wherein said aniline is 2,4,6-
2 trimethylaniline.

1 20. The method of claim 18, wherein said aniline is 2,6-
2 diisopropylaniline.

1 21. The method of claim 18, further comprising the step of
2 hydrogenating the nucleophilic carbene to produce a non-
3 aromatic nucleophilic carbene.

1 22. A method of performing ring closing metathesis, said
2 method comprising contacting a diterminal diene with a
3 catalytic complex under conditions appropriate, and for a
4 time sufficient to produce a cyclic alkene, wherein the
5 catalytic complex comprises:

6 a) a metal atom selected from the group
7 consisting of ruthenium or osmium;

8 b) at least one anionic ligand ligated to said
9 metal;


10 c) at least one nucleophilic carbene ligand ligated
11 to said metal;

12 d) a further ligand ligated to said metal; and

13 e) a carbon-containing ligand ligated to said metal,
14 wherein said carbon-containing ligand is
15 selected from the group consisting of


16 alkylidene, benzylidene, indenylidene, vinylidene,
17 and allenylidene.

1 23. The method of claim 22, wherein the catalytic complex
2 has the formula:



3
4 wherein M is Os or Ru;
5 C¹, C² and C³ are sp²-hybridized carbons, wherein
6 either or both of C¹ and C² are optionally absent;
7 R and R¹ are independently selected from the group
8 consisting of hydrogen, C₁-C₂₀ alkyl, C₂-C₂₀ alkenyl, C₂-C₂₀
9 alkynyl, C₂-C₂₀ alkoxy, carbonyl, aryl, C₁-C₂₀ carboxylate, C₁-
10 C₂₀ alkoxy, C₂-C₂₀ alkenyloxy, C₂-C₂₀ alkynyloxy, or aryloxy,
11 each R and R¹ optionally being substituted with C₁-C₅ alkyl,
12 halogen, C₁-C₆ alkoxy, or with a phenyl group substituted
13 with halogen, C₁-C₅ alkyl or C₁-C₅ alkoxy;
14 X and X¹ are independently selected from the group
15 consisting of anionic ligands; and
16 L and L¹ are selected from the group consisting of
17 nucleophilic carbenes, phosphine, sulfonated phosphine,
18 phosphite, phosphinite, phosphonite, ether, amine, amide,
19 sulfoxide, carbonyl, nitrosyl, pyridine and thioether,
20 wherein at least one of L or L¹ is a nucleophilic carbene.

1 24. The method of claim 23, wherein the nucleophilic
2 carbene has the formula:




3

4 wherein

5 Y and Y¹ are independently selected from the group
6 consisting of hydrogen, C₁-C₂₀ alkyl, C₂-C₂₀ alkenyl, C₂-C₂₀
7 alkynyl, C₂-C₂₀ alkoxy, C₂-C₂₀ alkoxy, C₂-C₂₀ alkenyloxy, C₂-C₂₀ alkyloxy, or aryloxy,
8 each Y and Y¹ optionally being substituted with C₁-C₅ alkyl,
9 halogen, C₁-C₆ alkoxy, or with a phenyl group substituted
10 with halogen, C₁-C₅ alkyl or C₁-C₅ alkoxy and;

12 Z and Z¹ are independently selected from the group
13 consisting of hydrogen, C₁-C₂₀ alkyl, C₂-C₂₀ alkenyl, C₂-C₂₀
14 alkynyl, C₂-C₂₀ alkoxy, C₂-C₂₀ alkoxy, C₂-C₂₀ alkenyloxy, C₂-C₂₀ alkyloxy, or aryloxy,
15 each Z and Z¹ optionally being substituted with C₁-C₅ alkyl,
16 halogen, C₁-C₆ alkoxy, or with a phenyl group substituted
17 with halogen, C₁-C₅ alkyl or C₁-C₅ alkoxy,
18 and wherein the ring can be optionally aromatic by
19 introduction of a further double bond in the ring.
20

1 25. The method of claim 22, wherein the catalytic complex
2 has the formula:



3

4

5

6 wherein

7 C¹, C² and C³ are sp²-hybridized carbons, wherein
8 either or both of C¹ and C² are optionally absent;

9 M is selected from the group consisting of Os and
10 Ru;


R and R¹ are independently selected from the group consisting of hydrogen, C₁-C₂₀ alkyl, C₂-C₂₀ alkenyl, C₂-C₂₀ alkynyl, C₂-C₂₀ alkoxycarbonyl, aryl, C₁-C₂₀ carboxylate, C₁-C₂₀ alkoxy, C₂-C₂₀ alkenyloxy, C₂-C₂₀ alkynyloxy, or aryloxy, each R and R¹ optionally being substituted with C₁-C₅ alkyl, halogen, C₁-C₆ alkoxy, or with a phenyl group substituted with halogen, C₁-C₅ alkyl or C₁-C₅ alkoxy;

X is an anionic ligand; and

L is a nucleophilic carbene; and

Ar is an aryl substituent, bonded to M by an η^6 bond.

26. The method of claim 25, wherein the nucleophilic carbene has the formula:



wherein

Y and Y¹ are independently selected from the group consisting of hydrogen, C₁-C₂₀ alkyl, C₂-C₂₀ alkenyl, C₂-C₂₀ alkynyl, C₂-C₂₀ alkoxycarbonyl, aryl, C₁-C₂₀ carboxylate, C₁-C₂₀ alkoxy, C₂-C₂₀ alkenyloxy, C₂-C₂₀ alkynyloxy, or aryloxy, each Y and Y¹ optionally being substituted with C₁-C₅ alkyl, halogen, C₁-C₆ alkoxy, or with a phenyl group substituted with halogen, C₁-C₅ alkyl or C₁-C₅ alkoxy and;

Z and Z¹ are independently selected from the group consisting of hydrogen, C₁-C₂₀ alkyl, C₂-C₂₀ alkenyl, C₂-C₂₀ alkynyl, C₂-C₂₀ alkoxycarbonyl, aryl, C₁-C₂₀ carboxylate, C₁-C₂₀ alkoxy, C₂-C₂₀ alkenyloxy, C₂-C₂₀ alkynyloxy, or aryloxy, each Z and Z¹ optionally being substituted with C₁-C₅ alkyl, halogen, C₁-C₆ alkoxy, or with a phenyl group substituted with halogen, C₁-C₅ alkyl or C₁-C₅ alkoxy,

19 and wherein the ring can be optionally aromatic by
20 introduction of a further double bond in the ring.

Abstract

1 Catalytic complexes including a metal atom having
2 anionic ligands, at least one nucleophilic carbene ligand,
3 and an alkylidene, vinylidene, or allenylidene ligand. The
4 complexes are highly stable to air, moisture and thermal
5 degradation. The complexes are designed to efficiently
6 carry out a variety of olefin metathesis reactions.

56656.N11




Fig. 1A




Fig. 1B




Fig. 1C




Fig. 2A




Fig. 2B




Fig. 2C




Fig. 3A




Fig. 3B




Fig. 3C




FIG. 4




FIG. 5

[illegible]

FIG. 6

COMBINED DECLARATION AND POWER OF ATTORNEY

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name,

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled CATALYST COMPLEX WITH CARBENE LIGAND, the specification of which

☒ is attached hereto.

☐ was filed on _____ as Application Serial No. _____ and was amended on _____.

☐ was described and claimed in PCT International Application No. _____ filed on _____ and as amended under PCT Article 19 on _____.

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose all information I know to be material to patentability in accordance with Title 37, Code of Federal Regulations, §1.56.

I hereby claim the benefit under Title 35, United States Code, §119(e)(1) of any United States provisional application(s) listed below:

U.S. SERIAL NO.	FILING DATE	STATUS
<u>60/099,722</u>	<u>September 10, 1998</u>	PENDING
<u>60/115,358</u>	<u>January 8, 1999</u>	PENDING

I hereby appoint the following attorneys and/or agents to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith: Scott C. Harris, Reg. No. 32,030; John R. Wetherell, Jr., Reg. No. 31,678; George E. Heibel, Reg. No. 42, 648; Richard P. Ferrara, Reg. No. 30,632; Stephan J. Filipek, Reg. No. 33,384; William J. Hone, Reg. No. 26,739; Stuart Macphail, Reg. No. 44,217; Greg O' Bradovich, Reg. No. 42,945; Andrew N. Parfomak, Reg. No. 32,431; John B. Pegram, Reg. No. 25,198; Frederick H. Rabin, Reg. No. 24,488; Gary A. Walpert, Reg. No. 26,098.

Address all telephone calls to Scott C. Harris at telephone number 619/678-5070.

Address all correspondence to Scott C. Harris, Fish & Richardson P.C., 4225 Executive Square, Ste. 1400, La Jolla, CA 92037.

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patents issued thereon.

COMBINED DECLARATION AND POWER OF ATTORNEY CONTINUED

Full Name of Inventor: Steven P. Nolan

Inventor's Signature: Sholan Date: Sept 8, 1999

Residence Address: 6196 Chatham Drive, Apt. 153, New Orleans, LA 70122

Citizen of: Canada

Post Office Address: 6196 Chatham Drive, Apt. 153, New Orleans, LA 70122

Full Name of Inventor: Jinkun Huang

Inventor's Signature: _____ Date: _____

Residence Address: 2000 Lakeshore Drive, Apt. D-208, New Orleans, LA 70148

Citizen of: China

Post Office Address: 2000 Lakeshore Drive, Apt. D-208, New Orleans, LA 70148

56932.N11